

**Salvin's albatross –**


**Bounty Is. population estimate**

**2013**

**Barry Baker, Katrina Jenz & Paul Sagar**

**Latitude 42 Environmental Consultants**

**NIWA**





# background

- endemic NZ species
- Bounty Islands  
98% global population (c.32 000 pairs) breeds on 7 islands
- Snares Western Chain c. 2,000 pairs

# WEST GROUP





# EAST GROUP



# CENTRE GROUP





## Project Aims

- develop population estimate for Salvin's albatross on the Bounty Islands for 2013

- estimate population trend in relation to previous aerial surveys & ground counts

# methods

- October 2010 & 2013 flew from Christchurch to the Bounty's
- fixed wing aircraft  
twin turboprop  
Cessna Conquest II
- population census at Bounty Is colonies using  
aerial photography
- no ground truthing 2010  
ground truthing 2013 Proclamation I.



# ground count protocol

- Proclamation I. 23/10/2013,  
same day as aerial photography
- to determine proportions of
  - nests containing eggs; &
  - non-breeding birds present in the colony


# ground count protocol

- 2 transects of variable length X 2m wide
- transect length determined by density of nesting birds, terminated when 100 active nests with eggs located.
- all nests with eggs 1 m either side of lines counted & spot marked; together with empty nests
- counts repeated every 2 hours, recording nests with eggs, empty nests with bird on, loafing birds, partners of incubating birds
- counts along transects undertaken by one observer at 1000, 1200, 1400 and 1600 hours


# photo counting protocol

- photomontages constructed using Adobe Photoshop software
- paintbrush tool mark off counted birds

# data assessment

A close-up photograph of a bird's wing and tail against a blue sky background. The bird's feathers are dark brown and grey, and its long, hooked beak is visible. The background is a soft, out-of-focus blue sky.

- all birds on the ground counted.
- each single bird assumed to represent an annual breeding pair.
- all images counted by one observer
- repeat counts of randomly selected montages by 2 other observers confirmed no evidence of observer bias in counting
- counts adjusted to reflect the proportion of non-breeding birds determined by ground-truthing in 2013



penguin

albatross

seal





# ground count results

Time	Transect #	Nest with eggs	Empty nests with bird on	Loafers	Total birds	Proportion birds breeding
10.00	1	100	10	28	138	0.72
	2	100	11	19	130	0.77
12.00	1	100	10	27	137	0.73
	2	100	14	19	133	0.75
14.00	1	100	11	29	140	0.71
	2	100	12	20	132	0.76
16.00	1	100	11	27	138	0.72
	2	100	11	19	130	0.77
<b>Mean</b>						<b>0.74</b>

# aerial counts

- total count October 2013 – mid incubation  
53,893 (CI 53,429 — 54,357)
- mean proportion breeding birds 0.74  
occupied nests containing eggs 0.90
- 39,995 (95% CI 39,595 - 40,395)  
annual breeding pairs in 2013
- most birds on Depot Is. 15,000

# Bounty Island raw counts & estimated annual breeding pairs

ISLAND	2010		2013	
	Raw	Corrected	Raw	Corrected
Molly Cap	3,361	2,494	4,390	3,258
Funnel I.	5,159	3,829	6,983	5,182
Spider I.	3,750	2,783	4,644	3,446
Depot I.	17,862	13,256	18,510	13,737
Penguin I.	2,203	1,635	1,407	1,044
Ruatara I.	5,313	3,943	6,754	5,012
Tunnel I.	2,333	1,731	4,629	3,435
Proclamation I.	2,851	2,116	6,576	4,880
<b>TOTAL</b>	<b>41,101</b>	<b>31,786</b>	<b>53,893</b>	<b>39,995</b>
	SE 203	232	178	200

## discussion points

- aerial survey is effective method of assessing population size of Salvin's albatross at the Bounty Islands
- proportion of non-breeding birds (25.8%) is high, but may be normal at this stage (mid-incubation period) of breeding
- developing annual correction factors in the absence of ground truthing not feasible from aerial photography (unlike white-capped albatross breeding in Auckland Is)
- non-breeder proportion likely to be much lower (c. 5%) 6 weeks earlier at end of egg laying period
- not appropriate to draw conclusions about population changes by combining aerial counts with earlier data sets
  - methods vary considerably
  - carried out at different times of breeding cycle

# Acknowledgements

Scientific and Technical Support:

Katrina Jenz, Graham Robertson, Matt Charteris,

Boat:

*Tiama*, Henk Haazan

Aircraft:

NZ Flying Doctor Service, Andrew Currie,  
Dion Currie, Grant Mitchell

DOC:

Pete McClelland, Brent Bevan, Jo Hiscock,  
Sharon Trainor, Igor Debksi, Ian Angus

Deepwater Group:

Richard Wells